

CEUTICACILLIN

W.S.P

(For vet. use only)

Compsition:

Each 1gm contains:

Amoxicillin trihydrate 700 mg
(Equivalent to 610 mg base amoxycillin)

Properties:

Amoxicillin is a semi-synthetic broad spectrum antibiotic from penicillin group. It's mainly time-dependant, bactericidal against Gram+ and Gram- bacteria, in particular Actinobacilluspleuropneumoniae, isolated in pigs and Escherichia coli isolated in broilers. Amoxicillin has been shown in vitro to be bactericidal against a wide range of gram-positive and gram-negative bacteria, which commonly cause disease in poultry ,and pigs such as - Actinobacillus spp. , Actinopyogenes, Erysipelothrixrhusiopathiae , Staphylococci , Streptococci , Pasteurellasp , Clostridium spp, Salmonella spp ,HaemophilusParagallinarum, and E.coli.

Target Species:

Poultry (Broiler chickens-Fattening turkey) and Pigs.

Indications:

Poultry(Broiler chickens-Fattening turkey):-

- Necrotic enteritis caused by clostridium perfringens.
- Pasteurellosis caused by Pasteurellamultocida.-
- Colibacillosis caused by E.coli.Pigs: Swine pleuropneumonia due to Actinobacilluspleuropneumoniae.

Dosage and adminstration:

Orally Via drinking water.

For Whole product:

Broiler chickens: 24.59 gm of product /1000 kg.b.wt. in drinking water /day/5 days (0.25 gm/ 1 liter).

Fattening turkeys: 24.59 gm-32.79 gm of product / 1000 kg.b.wt. in drinking water /day /5 days(0.35-0.25 gm/liter).

Pigs: 2 gm of product/ 61 kg.b.wt. in drinking water/day/4 days.

Warning and percuation:

- Not to be used simultaneously with neomycin since it blocks the absorption of oral penicillins.
- The uptake of medication by animals can be altered as a consequence of illness. In case of insufficient uptake of feed/ water animals should be treated parenterally.

Administration:

5 teiba st, From Alhouda wa Alnour st,
Mansoura, Dakahlia, Egypt.

factory:

Gamasa industrial zone 4 th area, Block B, No 6
Gamasa, Dakahlia, Egypt.

phone:

01273534552
050 272 96 96

email:

Sales@edpharma.com.eg
www.edpharma.com.eg

CEUTICACILLIN

W.S.P

(For vet. use only)

- Use of the product should be based on susceptibility testing of the bacteria isolated from animals. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria

Contra-indications:

- Do not give amoxicillin orally (or parenterally) to small herbivores such as guinea pigs, hamsters and rabbits as with other penicillins.
- Do not use in conjunction with bacteriostatics.
- Do not use to treat infections caused by Beta-lactamase producing organisms , as amoxicillin is ineffective.

Side effects:

Penicillins may cause hypersensitivity following administration.
Allergic reactions to penicillin may occasionally be serious.

Withdrawal time:

Meat:

Pigs: 6 days.

Broiler chickens: 1 day.

Turkeys for fattening: 5 days.

Its use is unauthorized in laying birds producing eggs for human consumption .
Do not use within 4 weeks from the start of laying.

Storage condations:

Store at a temperature not exceeding 30°C in a dry place used immediately after opening and reconstitution.

Packaging:

HDPE white plastic jar contains LDPE colorless plastic bag contains 20, 25, 50, 100, 250, 500, 1000, 2000, 5000, 10000 or 25000 gm water soluble powder and tightened with LDPE white plastic seal strip, The jar covered by HDPE white plastic cap with safety seal with outer label and inner leaflet.

**Manufactured by: The Arab Company for Gelatin And Pharmaceutical Products (Arab Caps)
For ED Pharma**

Administration:

5 teiba st, From Alhouda wa Alnour st,
Mansoura, Dakahlia, Egypt.

factory:

Gamasa industrial zone 4 th area, Block B, No 6
Gamasa, Dakahlia, Egypt.

phone:

01273534552
050 272 96 96

email:

Sales@edpharma.com.eg
www.edpharma.com.eg