

Doxyvestal

W.S.P

(For vet. use only)

Compostion:

Each 1 gm contains:

Doxycycline hyclate 1000 mg (Eq. To doxycycline base 867 mg)

Properties:

Doxycycline, a tetracycline derivative is effective against aerobic and anaerobic gram-positive and gram-negative bacteria, Rickettsia, Chlamydia, Mycoplasma, spirochetes and Protozoa.

Doxycycline inhibits bacterial protein synthesis intracellularly by binding on the 30-S ribosome subunits. This interferes with binding of aminoacetyl-tRNA to the acceptor site on the mRNA ribosome complex and prevents coupling of amino acids to the elongating peptide chains. The effect is bacteriostatic.

Indication:

Pigs: Treatment of the following specified infections of the respiratory tract caused by micro-organisms susceptible to doxycycline as:

- Atrophic rhinitis caused by Pasteurella multocida and Bordetella bronchiseptica;
- Bronchopneumonia caused by Pasteurella multocida, Streptococcus suis and Mycoplasma hyorhinis;
- Pleuropneumonia caused by Actinobacillus pleuropneumoniae.

Broiler Chickens:

- Infections of the respiratory tract caused by Mycoplasma spp., Escherichia coli, Haemophilus, paragallinarum and Bordetella avium;
- Enteritis caused by Clostridium perfringens and Clostridium colinum.

Target species:

Pig and Broiler chickens.

Dosage and administration:

Route Orally via drinking water.

Dose of whole product:

Pig: 0.75 g of the product equivalent to one day of treatment of 50 kg pig for 5 days.

Broiler Chickens: 0.25 g of the product / liter for 5-3 consecutive days.

Administration:

5 teiba st, From Alhouda wa Alnour st,
Mansoura, Dakahlia, Egypt.

factory:

Gamasa industrial zone 4 th area, Block B, No 6
Gamasa, Dakahlia, Egypt.

phone:

01273534552
050 272 96 96

email:

Sales@edpharma.com.eg
www.edpharma.com.eg



Doxyvestal

W.S.P

(For vet. use only)

Pre-ruminating calves: 1 gm of the product /12.5 kg bw twice daily, for 7 to 14 days.

Warning & Precautions:

Use of the product should be based on susceptibility testing.

Absorption of doxycycline can be decreased in presence of high quantities of calcium, Iron , mg in diet.

Contra – Indications:

- Do not use in animals previously demonstrated hypersensitivity to tetracyclines.
- Not permitted for use in laying birds producing eggs for human consumption.
- Do not use in animals suffering from kidney problems.

Adverse Effects:

As for all tetracyclines, allergic reactions or photo sensitization occur.

Yellowing of bones and teeth and Enamel hypoplasia may occur after use in young animals.

Withdrawal time:

In pig (Meat and offal): 8 days.

Broiler Chickens: 10 days

Storage conditions:

Store at temperature not more than 30°C. in dry place and used immediately after opening and reconstitution.

Packaging:

White plastic (HDPE) Jar containing 10 gm & 1000 gm powder with an outer label sealed with aluminum seal lined inside by (HDPE) plastic film, closed by white plastic (HDPE) cap, with (LDPE) plastic foam imbedded in cap.

**Manufactured by Amcomed For Pharmaceutical And Chemical Industries
for Ed pharma**

Administration:

5 teiba st, From Alhouda wa Alnour st,
Mansoura, Dakahlia, Egypt.

factory:

Gamasa industrial zone 4 th area, Block B, No 6
Gamasa, Dakahlia, Egypt.

phone:

01273534552
050 272 96 96

email:

Sales@edpharma.com.eg
www.edpharma.com.eg