

# ECOMARBOFLOX

Solution for Injection  
(For vet. use only)

## Composition:

### Each 1 ml contains:

Marbofloxacin                      160mg.

## Properties:

Marbofloxacin is a synthetic, bactericidal antimicrobial, belonging to the fluoroquinolone group. It acts by inhibition of DNA gyrase enzyme of microbial cell. The in vitro activity of marbofloxacin has been demonstrated towards *Pasteurella multocida*, *Mannheimia haemolytica* and *Escherichia coli*.

## Target species:

Cattle.

## Indications:

- Therapeutic treatment of respiratory infections caused by sensitive strains of *Pasteurella multocida* and *Mannheimia haemolytica*.
- Acute mastitis in lactating cows caused by *E. coli*.

## Dosage and Administration:

**Route:** by single IM or IV injection.

### Dose:

10 ml of product/160 kg.b.wt.

## Warnings and Precautions:

Use of the product should only be based on susceptibility testing.

- If the volume to be injected intramuscularly is more than 20 ml, it should be divided between two or more injection sites.
- To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing.

## Contraindications:

- Do not use in animals with known hypersensitivity to fluoroquinolones or to any of the excipients.
- Do not use in cases where the pathogen involved is resistant to other fluoroquinolones (cross resistance).

### Administration:

5 teiba st, From Alhouda wa Alnour st,  
Mansoura, Dakahlia, Egypt.

### factory:

Gamasa industrial zone 4 th area, Block B, No 6  
Gamasa, Dakahlia, Egypt.

### phone:

01273534552  
050 272 96 96

### email:

Sales@edpharma.com.eg  
www.edpharma.com.eg

# ECOMARBOFLOX

Solution for Injection  
(For vet. use only)

## Side effects:

- Administration by the intramuscular route may cause rare transient local reactions such as pain and swelling at the injection site which may persist up to 7 days after injection.
- Fluoroquinolones are known to induce arthropathies.
- Lesions of the joint cartilage were observed in some animals treated at 10 mg/kg or 30 mg/kg for three times the recommended treatment duration, but did not induce clinical signs. Moreover, no sign of overdose was observed throughout this study

## Withdrawal Time:

**Meat and offal:** 6 days.

**Milk:** 48 hours.

## Storage Conditions:

Store at temperature not exceeding 30°C and to be used immediately after opening.

## Packaging:

Amber glass (type II) vial contains 100 or 250 ml injectable solution and closed with a bromobutyl rubber stopper over sealed with an aluminum.

**Manufactured by: Delta Grand pharma for pharmaceutical industries  
For Ed Pharma**

### Administration:

5 teiba st, From Alhouda wa Alnour st,  
Mansoura, Dakahlia, Egypt.

### factory:

Gamasa industrial zone 4 th area, Block B, No 6  
Gamasa, Dakahlia, Egypt.

### phone:

01273534552  
050 272 96 96

### email:

Sales@edpharma.com.eg  
www.edpharma.com.eg