

Composition

Contain per 1 ml :

Active ingredients:

Amprolium hydrochloride	240 mg.
equivalent to amprolium base	212.2 mg.

Properties:

Amprolium is a coccidiostat belonging to the thiamine (Vitamin B1) analogue group. It acts as a competitive antagonist of thiamine transport mechanisms. Amprolium affects coccidia by competing with thiamine in their metabolic enzyme system, thus interfering with metabolism of carbohydrates necessary for multiplication or survival of coccidian. Amprolium is active against *E. tenella*, *E. necatrix*, *E. acervulina*, *E. Maxima*, *E. Brunette*, *E. adenoides*, *E. gallapavonis* and *E. meleagrimitis*.

Indications:

Broiler chickens:

- Treatment of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima* and *E. brunetti*.

Turkeys:

- Treatment of coccidiosis caused by *E. adenoides*, *E. gallapavonis* and *E. meleagrimitis*.

Target species:

Broiler chickens and turkeys.

Dosage and administration:

Route of administration: Orally via drinking water.

Dose of whole product:

Broilers chickens and turkeys:

942.5 mg of the product / 1 liter drinking water daily, for 7-5 days.

Warnings & Precautions:

- Frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.
- The product is not intended for a preventive use.
- The product may cause airway, eye and skin irritation.
- Wear impervious gloves and protective glasses when handling the product.
- Avoid inhalation of vapours.
- Do not eat, drink or smoke when handling the product.
- People with known hypersensitivity to amprolium or excipients should avoid contact with the product.

Administration:

5 teiba st, From Alhouda wa Alhour st,
Mansoura, Dakahlia, Egypt.

factory:

Gamasa industrial zone 4 th area, Block B, No 6
Gamasa, Dakahlia, Egypt.

phone:

01273534552
050 272 96 96

email:

Sales@edpharma.com.eg
www.edpharma.com.eg

ZOARIUM

- Avoid contact with the skin and eyes
- In the case of contact with skin or eyes, wash the affected area with clean running water immediately and remove any contaminated clothes.
- Wash hands and exposed skin after use.

Contraindications:

- Do not use in cases of known hypersensitivity to the active substance or any of the excipients.
- Do not use simultaneously with products containing vitamin B-complex.

Adverse reactions:

Prolonged uses can produce thiamine deficiencies.

Withdrawal times:

Chickens and Turkeys: Meat and offal: Zero days.

Eggs: Zero days.

Storage conditions:

Store at temperature not exceeding 30°C, in dry place and used immediately after opening and reconstitution.

Packaging:

Opaque HDPE plastic jar contains LDPE colorless plastic bag contains 100, 250, 500, 1000 g powder and tightened with LDPE plastic seal strip, The jar covered by HDPE opaque plastic cap with HDPE safety seal with outer label.

Administration:

5 teiba st, From Alhouda wa Alnour st,
Mansoura, Dakahlia, Egypt.

factory:

Gamasa industrial zone 4 th area, Block B, No 6
Gamasa, Dakahlia, Egypt.

phone:

01273534552
050 272 96 96

email:

Sales@edpharma.com.eg
www.edpharma.com.eg